
multitemperature data and diffuse scattering in X-ray crystallography



James S Fraser UCSF







Daniel Keedy



Robert Thorne (Cornell)



Henry van den Bedem (SLAC)



Andrew VanBenschoten





Michael Wall (LANL)

Nick Sauter, Aaron Brewster, Paul Adams, Pavel Afonine (LBNL), Tom Terwilliger (LANL), Sacha Urzhumtsev (CNRS)

XFELs and structural biology: It's about time!


We are transitioning from static structural biology....






...to dynamic structural biology


How are **allosteric** perturbations communicated intramolecularly to alter protein function?




Proteins often populate multiple conformations in crystals




Conformational heterogeneity can be **static** or **dynamic**




Conformational heterogeneity can be **static** or **dynamic**




Temperature can **shift** the relative populations of confirmations in the crystal



Temperature can **shift** the relative populations of confirmations in the crystal



Temperature can **shift** the relative populations of confirmations in the crystal



Hypotheses: (1) shifting temperature exposes conformations near the "ground" state; (2) these new conformations are used by the protein in physiological mechanisms

Conformational dynamics are at the core of three critical problems in biology

We want to:

design macromolecules with new (unnatural) functions

understand how mutations alter protein function in disease

discover small molecules drugs to modulate protein function

Conformational dynamics are at the core of three critical problems in biology

We want to:


design macromolecules with new (unnatural) functions


understand how mutations alter protein function in disease

discover small molecules drugs to modulate protein function

Hypotheses: (1) shifting temperature exposes conformations near the "ground" state; (2) these new conformations are used by the protein in physiological mechanisms

Conventional X-ray experiments are performed at cryogenic temperatures







...and data collection strategies to minimize radiation damage

Fraser...Alber, PNAS, 2011

now: XFELs!

Cryocooling contracts the lattice - and protein







Juers and Matthews, JMB, 2002

Frauenfelder et al, Biochemistry, 1987

Cryocooling causes **local changes** in structure and conformational heterogeneity






Temperature-dependent X-ray diffraction as a probe of protein structural dynamics


Hans Frauenfelder, Gregory A. Petsko* & Demetrius Tsernoglou

Department of Physics, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801 and

Department of Biochemistry, Wayne State University School of Medicine, Detroit, Michigan 48201

Cryocooling **slowly** shrinks proteins, remodels side chains, and improves packing






Room temperature data are more consistent than cryocooled data




Keedy...Fraser, Structure, 2014

The challenge is to deconvolute multiple conformations from an **ensemble averaged** density map



Ringer is an assay for electron density




0.3 and 1σ


Terry Lang Tom Alber

UC Berkeley


Side chains are built into high electron density regions




0.3 and 1σ




Only 5% of residues are modeled in multiple conformations






...but electron density sampling reveals multiple peaks


Ringer peaks are rotameric (non-random)



Most peaks represent hidden conformations




0.3 and 1σ




 $\chi 1 - \chi 2$ correlation plot

~600 structures at 1.5Å or better

Ringer gives us confidence in signals at low electron density









Terry Lang Tom Alber

UC Berkeley




James Holton

LBL/UCSF

RINGER:

Lang, Ng...Alber *Protein Science*, 2010

END/RAPID: Lang, Holton, Fraser, Alber PNAS, 2014



qFit automates
multiconformer
side chain and
backbone
modeling




Henry van den Bedem (SSRL)



Daniel Keedy

Alternative conformations (refined occupancies) model anharmonic motions better than B-factors








Henry van den Bedem (SSRL)



Daniel Keedy

Alternative conformations (refined occupancies) model anharmonic motions better than B-factors



mFo-DFc 2mFo-DFc -3.0σ 1.0σ $+3.0\sigma$ 0.5σ




Henry van den Bedem (SSRL)



Daniel Keedy

van den Bedem...Deacon, Acta D, 2009; Keedy, Fraser, van den Bedem, PLOS CompBiol, 2015

~35% of side chain conformations are changed by cryocooling




29% of all buried residues


Fraser...Alber, *PNAS*, 2011 with Henry van den Bedem (JCSG)

Coupled side chain motions explain NMR relaxation, enabling design of mutations to demonstrate

dynamics-function link





Conformational differences between room temperature and cryogenic temperatures using SFX at **XFELs**




Liu...Cherezov, Science, 2013

Conformational dynamics are at the core of three critical problems in biology







Can temperature perturbation expose the protein conformations

used by design, disease, and drugs?

We want to:


design macromolecules with new (unnatural) functions


understand how mutations alter protein function in disease

discover small molecules drugs to modulate protein function

Hypotheses: (1) shifting temperature exposes conformations near the "ground" state; (2) these new conformations are used by the protein in physiological mechanisms

Designed proteins (e.g. Ubiquitin variants to inhibit USP7) are currently **too dynamic**











After directed evolution, dynamics are reduced....



How **different** are dynamics of designed, artificially evolved, and natural proteins?



We want to:


design macromolecules with new (unnatural) functions

understand how mutations alter protein function in disease

discover small molecules drugs to modulate protein function

Hypotheses: (1) shifting temperature exposes conformations near the "ground" state; (2) these new conformations are used by the protein in physiological mechanisms


Why are these two resistance mutations in TEM-1 incompatible?



+/- inhibitor

Different resistance mutants cannot be combined

due to dynamics



suggests a method discover combination therapies with huge barriers to resistance





Eynat Dellus-Gur, Dan Tawfik (Weizmann)

Dellus-Gur...Fraser, Tawfik, JMB, 2015

We want to:

design macromolecules with new (unnatural) functions


understand how mutations alter protein function in disease

discover small molecules drugs to modulate protein function


Hypotheses: (1) shifting temperature exposes conformations near the "ground" state; (2) these new conformations are used by the protein in physiological mechanisms


Cryptic allosteric fragment sites are quenched by cryocooling

Active site heterogeneity differs between cryo and room temperature



Cryptic site:






changes small molecules discovered by in silico docking procedures

Fischer, Coleman, Fraser, Shoichet Nature Chemistry, 2014


Fischer, Shoichet, Fraser *ChemBioChem*, 2015





Marcus Fischer, Brian Shiochet (UCSF)

M2 Channel water density is temperature sensitive - ordered around drug binding sites and **pH sensitive**



Thomaston...Woldeyes, Fraser...DeGrado, PNAS, 2015




Jessica Thomaston (DeGrado Lab)




Rahel Woldeyes

Low-occupancy features present at room temperature are dynamically accessed conformations and can provide new mechanistic insights




Temperature can **shift** the relative populations of confirmations in the crystal



Does this happen collectively? How complex is the energy landscape?

CypA conformational heterogeneity is sensitive to

temperature changes





Daniel Keedy






Matt Warkentin, Robert Thorne (Cornell)

CypA conformational heterogeneity is sensitive to

temperature changes






Daniel Keedy







Matt Warkentin, Robert Thorne (Cornell)



Different regions of the protein have their disorder arrested at **different temperatures**







(one reason why it has been challenging for us to **extrapolate** from 100K to 293K X-ray Data)

Is the response to temperature correlated across the CypA "dynamic network"?


CypA temperature response does not favor a single **coupled transition**





NMR relaxation dispersion experiments may sense population shuffle upon Phe113 rotameric switch





Population shuffle: alternative conformations do not segregate neatly into A/B - through "clash-relief" pathways




































Residues linked together in **CONTACT** networks






Sensitive to: ligands, mutations, temperature Predictive of: chemical shift perturbations, collective motions

van den Bedem...Fraser, *Nature Methods*, 2013

Visualizing networks of mobility in proteins NEWS AND VIEWS


Mark A Wilson


NATURE METHODS | VOL.10 NO.9 | SEPTEMBER 2013 | 835




Happily for future methods developers, there is an opportunity to improve the modeling of mobility in crystallographic data. Out of necessity, van den Bedem et al. 1 used a stringent steric overlap criterion for determining which residues are correlated. However, more subtly correlated motions will be missed with this approach and are difficult to experimentally characterize. A possible solution to this is the use of X-ray diffuse scattering data, which contain information about all correlated motions in the crystal. The value of diffuse scattering data has been known for decades^{8,9}, but, owing to multiple technical challenges, these data are still routinely discarded¹⁰. Future improvements in this direction will allow the full extent of X-ray diffraction from crystals to be collected and analyzed, thus providing a fully integrated view of the connection between protein structure and correlated motions.

Diffuse scattering can **distinguish** different models of coupled heterogeneity





Small features - between unit cells | Large features - within unit cells







Fig. 1 X-ray diffraction data from insulin. a, A 40-h exposure of a stationary crystal (3-fold axis nearly vertical) recorded with focused, monochromatic Cu K_{a_1} radiation from an Elliott rotating anode. The data (corrected for camera background and polarization) are displayed out to radius $R = 0.45 \text{ Å}^{-1}$. Bragg reflections are overexposed. The sharp arcs are due to diffraction from the Al foil window of the He beam tunnel. The colour table (optical density range 0-2 OD units) was constructed to distinguish small variations in intensity up to 0.5 OD units. b, Bragg reflections and haloes digitally separated by subtracting the smoothly-varying diffuse scattering component from the film data. The inset wedge shows the estimated circularly symmetrical Compton-plus water scattering. c, Variational scattering evaluated from the difference between a and the two components in b. The colour table scale in b and c is $1.5 \times$ that in a. Each intensity step in c equals 0.02 OD units in the data.

Caspar et al, Nature, 1988


Welberry et al, Acta B, 2011

Given an ensemble, **Guinier's equation** relates large scale features (along hkl) to conformations




Van Benschoten...Fraser, Acta D, 2015


Given an ensemble, **Guinier's equation** relates large scale features (along hkl) to conformations



Van Benschoten...Fraser, Acta D, 2015


Long (microsecond) molecular dynamics simulations of crystals provide mechanisms for diffuse scattering







Wall et al, PNAS, 2014


Agreement of anisotropic features is currently moderate, improvements needed for sampling and scoring?







New data sets are needed to advance diffuse scattering










Wall, Ealick, and Gruner, PNAS 1997




Andrew VanBenschoten




Michael Wall (LANL)





TLS models are poor at explaining diffuse intensities, but **liquid like motions** are better






Conformational dynamics are at the core of three critical problems in biology








Temperature perturbations expose the *coupled* protein conformations used by design, disease, and drugs

"UHH ..." Saul Cortez



Jennifer Li






David Mavor


BEER

OR






Dr. Lin Liu












Dr. Daniel Keedy





Ben Barad





Rahel Woldeyes



ANSWERS THE

Dr. Mike **Thompson**



OBSESSED

Erin Poss



Alex Wolff


Pooja Suresh

1/2









Justin

Biel





